



How to reduce anterior resection syndrome and post-operative complication after rectosigmoid resection

Giulio Ricotta ¹, Elodie Gauroy¹, Anne-Sophie Navarro ,¹ Alejandra Martinez ,^{1,2} Gwenael Ferron ^{1,3}

¹Department of Surgical Oncology, Institut Universitaire du Cancer Toulouse Oncopole
 Departement de chirurgie, Toulouse, Languedoc-Roussillon Midi, France

²Department of Surgical Oncology and INSERM CRCT Team 1, Institut Universitaire du Cancer Toulouse Oncopole, Toulouse, Occitanie, France

³Department of Surgical Oncology and INSERM CRCT Team 19, Oncogenesis of Sarcomas, Institut Universitaire du Cancer Toulouse Oncopole, Toulouse, Occitanie, France

Correspondence to

Dr Giulio Ricotta, Department of Surgical Oncology, Institut Universitaire du Cancer Toulouse Oncopole
 Departement de chirurgie, Toulouse, Languedoc-Roussillon Midi, France;
 drgricotta@gmail.com

Accepted 18 March 2024



© IGCS and ESGO 2024. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Ricotta G, Gauroy E, Navarro A-S, et al. *Int J Gynecol Cancer* Published Online First: [please include Day Month Year]. doi:10.1136/ijgc-2024-005306

SUMMARY

Intestinal surgery is often required in the management of patients with gynecological malignancies, and the rectosigmoid colon is the bowel segment most frequently involved.^{1,2}

In rectal cancer, the total mesorectal excision technique represents the standard procedure for surgical excision. However, since its introduction, the risk of anastomotic leakage and pelvic infection has increased. Moreover, total mesorectal excision is associated with pelvic and rectal autonomic nerve injury, which may cause 'anterior resection syndrome' (defined as disordered bowel function after rectal resection), leading to a detriment in quality of life.³



Figure 1 Photo showing the close rectal dissection

In gynecological malignancy, in cases where there is no deep macroscopic mesorectal disease, total mesorectal excision is not essential. Close rectal dissection technique is based on mesorectal

INTERNATIONAL JOURNAL OF
GYNECOLOGICAL CANCER

How to reduce anterior resection syndrome and post-operative complication after recto-sigmoid resection

Giulio Ricotta¹, MD; Elodie Gauroy¹, MD; Anne Sophie Navarro¹, MD; Alejandra Martinez^{1,2}, MD PhD; Gwenaël Ferron^{1,3}, MD PhD



 @IJGOnline

Copyright © 2020 BMJ Publishing Group Ltd, International Gynecologic Cancer Society, & European Society of Gynaecological Oncology. All rights reserved.

Video 1 Close rectal dissection technique

Video article

sparing, and allows the preservation of the superior rectal artery while minimizing autonomic nerve damage.

In a review evaluating these two different surgical procedures in rectal benign disease, close rectal dissection was associated with reduced nerve injury and pelvic sepsis, with a lower rate of anastomotic leakage and improved bowel function.⁴ These results were confirmed by Son et al in a retrospective study on patients with ovarian cancer, which showed no differences in oncologic outcomes.²

In this video, we present an anterior pelvic exenteration associated with a rectosigmoid resection with the close rectal dissection technique in a woman aged in her 50s, who had been treated with definitive chemoradiotherapy and brachytherapy for locally advanced cervical cancer. Ten months after the end of treatment, the patient locally recurred, and an anterior pelvic exenteration (type I Magrina) with a rectosigmoid resection with close rectal dissection was performed.

We describe the procedure of rectosigmoid resection with the close rectal dissection technique. To summarize, rectosigmoid resection is often necessary to achieve complete cytoreduction in gynecological malignancies. In cases of no macroscopic deep mesorectal localization of disease and without massive Douglas pouch involvement, total mesorectal excision is unnecessary. Close rectal dissection allows better anastomotic vascularization by preserving the superior rectal artery and reducing nerve injury, with a lower rate of anastomotic leakage, pelvic sepsis, and improved bowel function. Unfortunately, this surgical technique is not widely applied by surgeons, even when technically feasible. With this video, we aim to standardize its use (Video 1).

Twitter Giulio Ricotta @GiRicotta and Alejandra Martinez @Alejandra

Contributors GR, EG, ASN: conceptualization, video editing, and writing-original draft. AM: conceptualization, project administration, supervision, and writing-

review. GF: conceptualization, project administration, surgery and video recording, supervision, and writing-review. GR: guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement There are no data in this work.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Giulio Ricotta <http://orcid.org/0000-0001-5897-5650>

Anne-Sophie Navarro <http://orcid.org/0000-0003-3328-7055>

Alejandra Martinez <http://orcid.org/0000-0002-7633-3536>

Gwenael Ferron <http://orcid.org/0000-0002-8545-4700>

REFERENCES

- 1 Rubin SC, Benjamin I, Hoskins WJ, et al. Intestinal surgery in gynecologic oncology. *Gynecol Oncol* 1989;34:30–3.
- 2 Son J-H, Kim J, Shim J, et al. Comparison of posterior Rectal dissection techniques during Rectosigmoid colon resection as part of Cytoreductive surgery in patients with epithelial ovarian cancer: close Rectal dissection versus total Mesorectal Excision. *Gynecol Oncol* 2019;153:362–7.
- 3 Bryant CLC, Lunniss PJ, Knowles CH, et al. Anterior resection syndrome. *Lancet Oncol* 2012;13:e403–8.
- 4 Nally DM, Kavanagh DO, Winter DC. Close Rectal dissection in benign diseases of the Rectum: A review. *Surgeon* 2019;17:119–26.