

Variables	Risk factors	HR (95% CI)	<i>p</i>
<i>Locoregional recurrence</i>			
Age		1.00 (0.94–1.06)	0.94
Tumour size	>4 cm vs. ≤4 cm	2.83 (1.45–5.50)	0.002
Histology	Non-end vs. end.	1.76 (0.85–3.62)	0.13
Grade	III vs. I-II	1.67 (0.80–3.51)	0.17
LVSI	Absent vs. present	0.73 (0.37–1.45)	0.37
Cervical glandular inv	Absent vs. present	0.78 (0.36–1.67)	0.52
Cervical stromal inv	Absent vs. present	1.69 (0.75–3.81)	0.21
Ln metastasis	Present vs. absent	1.21 (0.44–3.32)	0.72
Stage	III vs. I-II	3.34 (1.08–10.30)	0.04
<i>Distant metastasis</i>			
Age		1.02 (0.99–1.06)	0.2
Tumour size	>4 cm vs. ≤4 cm	1.42 (0.97–2.08)	0.08
Histology	Non-end vs. end.	1.47 (0.95–2.26)	0.08
Grade	III vs. I-II	1.66 (1.07–2.57)	0.02
LVSI	Absent vs. present	0.88 (0.58–1.33)	0.53
MI	≥1/2 vs. <1/2	1.44 (0.67–3.11)	0.35
Cervical glandular inv	Absent vs. present	0.60 (0.37–0.98)	0.04
Cervical stromal inv	Absent vs. present	0.44 (0.27–0.72)	0.001

Ln metastasis	Present vs. absent	2.85 (1.35–6.00)	0.006
Stage	III vs. I-II	2.17 (0.94–5.02)	0.07