

Table S2. Comparison of patient demographics and outcomes by mode of presentation for women with high grade serous ovarian cancer

	Fast-track pathway N=119	Emergency N=7	Other outpatients N=27
Age, years* mean (S.D), p=0.031	65.0 (10.1)	55 (16.1)	66.7 (10.2)
Stage** n (%), p=0.459			
Early stage			
1	12 (10.1)	2 (28.6)	2 (7.4)
2	18 (15.1)	0 (0.0)	4 (14.8)
Late stage			
3A	16 (13.5)	1 (14.3)	1 (3.7)
3B	11 (9.2)	0 (0.0)	0 (0.0)
3C	48 (40.3)	1 (14.3)	15 (55.6)
4A	7 (5.9)	0 (0.0)	1 (3.7)
4B	4 (3.4)	2 (28.6)	2 (7.4)
Not available	3 (2.5)	1 (14.3)	2 (7.4)
Performance status** n (%), P=0.611			
0	90 (75.6)	4 (57.1)	22 (81.5)
1	22 (18.5)	3 (42.9)	2 (7.4)
2	6 (5.0)	0 (0.0)	1 (3.7)
3	0 (0.0)	0 (0.0)	1 (3.7)
Not available	1 (0.9)	0 (0.0)	1 (3.7)
Cytoreduction rate** n (%), P=0.920			
Complete	73 (61.3)	4 (57.1)	18 (66.7)
Residual <1cm	18 (15.1)	1 (14.3)	3 (11.1)
Residual ≥1cm	2 (1.7)	0 (0.0)	2 (7.4)
Inoperable	9 (7.6)	0 (0.0)	2 (7.4)
Not available	17 (14.3)	2 (28.6)	2 (7.4)

* The ANOVA one-way test was used to calculate the p-value

**The Kruskal-Wallis H test was used to calculate the p-value